

MEDICATION GUIDE
RASONQUE™ (RAS-ON-cue)
(daraxonrasib)
tablets

What is the most important information I should know about RASONQUE?

RASONQUE can cause serious side effects, including:

- **Skin reactions.** Skin reactions may be severe. To help reduce your risk of skin reactions and protect your skin when starting and during treatment with RASONQUE:
 - use a topical steroid cream on your face and chest
 - use a moisturizing cream on your skin
 - limit your time in the sun and use sunscreen (broad spectrum SPF 30 or higher) and other sun protection measures such as clothing and hats if you must be in the sun
 - take any other medicines if prescribed by your healthcare providerTell your healthcare provider right away if you develop any signs or symptoms of skin reactions, including:
 - rash
 - itching
 - red, swollen, and painful skin around your finger or toenails
 - dry skin
 - cracked skin
- **Swelling or sores in the mouth (stomatitis).** Mouth sores may be severe. Your healthcare provider may prescribe a mouthwash or other medicines to treat your mouth reactions. Tell your healthcare provider right away if you develop trouble eating, talking, swallowing, or develop any new or worsening signs or symptoms in your mouth, including:
 - pain
 - redness
 - swelling
 - ulcers or sores
- **Diarrhea.** Diarrhea may be severe. If you develop diarrhea during treatment with RASONQUE, your healthcare provider may prescribe medicines to treat your diarrhea. Tell your healthcare provider right away if you develop new or worsening diarrhea or if the number of bowel movements you have in a day increases by 6 or more.

See “**What are the possible side effects of RASONQUE?**” for more information about side effects.

What is RASONQUE?

RASONQUE is a prescription medicine used to treat adults with a type of pancreatic cancer called pancreatic adenocarcinoma:

- that has spread to other parts of the body, **and**
- who have received at least 1 prior treatment or who cannot receive a combination of cancer treatments.

It is not known if RASONQUE is safe and effective in children.

Before taking RASONQUE, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. RASONQUE can harm your unborn baby.
Females who are able to become pregnant:
 - Your healthcare provider will do a pregnancy test before you start treatment with RASONQUE.
 - Use effective birth control (contraception) during treatment with RASONQUE and for 1 week after the last dose.
 - Tell your healthcare provider right away if you become pregnant during treatment with RASONQUE.**Males with female partners who are able to become pregnant:**
 - Use effective contraception during treatment with RASONQUE and for 1 week after the last dose.
- are breastfeeding or plan to breastfeed. It is not known if RASONQUE passes into your breast milk. **Do not** breastfeed during treatment and for 1 week after your last dose of RASONQUE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. RASONQUE can affect the way other medicines work, and other medicines can affect how RASONQUE works, and may increase your risk of side effects.

How should I take RASONQUE?

- Take RASONQUE exactly as your healthcare provider tells you to take it. Do not change your dose or stop taking RASONQUE unless your healthcare provider tells you to.
- Take RASONQUE 1 time each day, at the same time each day.
- Take RASONQUE with or without food.
- Swallow RASONQUE tablets whole. **Do not** chew, crush, or split tablets.
- If you miss a dose, take the dose as soon as you remember. If it has been more than 4 hours, skip the missed dose and take your next dose at your next regularly scheduled time. **Do not** take 2 doses at the same time to make up for a missed dose.
- If you vomit after taking a dose, do not take an extra dose. Take your next dose at your regularly scheduled time.

What are the possible side effects of RASONQUE?**RASONQUE can cause serious side effects, including:**

- See “What is the most important information I should know about RASONQUE?”
- **Tears in your stomach or intestines (gastrointestinal perforation).** Gastrointestinal perforation may be severe and lead to death. Tell your healthcare provider right away if you develop any of the following signs or symptoms:
 - tenderness or pain in your stomach area (abdomen)
 - blood in the stool or black stool that looks like tar that is severe or does not go away
 - nausea, vomiting, or vomiting blood
 - fever or chills
- **Lung or breathing problems (pneumonitis).** Lung or breathing problems may be severe and lead to death. Tell your healthcare provider right away if you develop any new or worsening breathing problems, including:
 - cough
 - shortness of breath
 - trouble breathing
 - wheezing

The most common side effects of RASONQUE include:

- | | |
|----------------------------------|---------------------------------|
| • rash | • vomiting |
| • diarrhea | • stomach (abdominal) pain |
| • swelling or sores in the mouth | • swelling in the arms and legs |
| • nausea | • decreased appetite |
| • tiredness or weakness | • bleeding |

Certain abnormal blood tests are common during treatment with RASONQUE. Your healthcare provider will monitor you for abnormal blood tests and treat you if needed.

Your healthcare provider may decrease your dose, or temporarily or permanently stop treatment with RASONQUE if you develop certain side effects.

These are not all of the possible side effects of RASONQUE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store RASONQUE?

Store RASONQUE at room temperature between 68°F to 77°F (20°C to 25°C).

Keep RASONQUE and all medicines out of the reach of children.

General information about the safe and effective use of RASONQUE.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use RASONQUE for a condition for which it was not prescribed. Do not give RASONQUE to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about RASONQUE that is written for health professionals.

What are the ingredients in RASONQUE?

Active ingredient: daraxonrasib

Inactive ingredients: butylated hydroxytoluene, colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl cellulose, hydroxypropyl methylcellulose acetate succinate, magnesium stearate, mannitol, and microcrystalline cellulose. The tablet film coating contains FD&C blue #2/indigo carmine aluminum lake, glyceryl mono and dicaprylocaprate, polyvinyl alcohol, sodium lauryl sulfate, talc, and titanium dioxide.

Manufactured for: Revolution Medicines, Inc. Redwood City, CA 94063 USA

© 2026 Revolution Medicines, Inc.

For more information, go to www.RASONQUE.com or call 1-844-2-REVMED (1-844-273-8633).